

Table II. *Effects of different levels of sumac seed powder (SSP) on growth performance of broiler chickens.*

Item	Control	0.05% SSP	0.10% SSP	0.15% SSP	0.20% SSP	SEM	P-value
Day 1-21							
Body weight (g/bird) at day 21	715.8	697.8	682.1	712.5	683.1	15.84	0.450
Average feed intake (g)	1,002.3	974.0	958.6	996.0	962.3	20.46	0.481
Feed conversion ratio (FCR)	1.40	1.39	1.40	1.39	1.40	0.008	0.890
Day 21-42							
Body weight (g/bird) at day 42	1,241.3	1,061.0	1,208.3	1,072.0	1,178.6	97.30	0.602
Average feed intake (g)	2,512.6	2,162.3	2,418.0	2,254.3	2,379.6	170.00	0.634
Feed conversion ratio (FCR)	2.02	2.04	2.01	2.10	2.02	0.037	0.461
Day 1-42							
Body weight (g/bird)	1,957.1	1,758.8	1,890.5	1,784.50	1,861.8	90.6	0.561
Average feed intake (g)	3,515.0	3,136.3	3,376.6	3,250.3	3,342.6	160.6	0.563
Feed conversion ratio (FCR)	1.79	1.78	1.78	1.82	1.79	0.016	0.522